

About the Matrix model computation of three-graviton scattering

Marco Fabbrichesi, Gabriele Ferretti and Roberto Iengo

*Istituto Nazionale di Fisica Nucleare, Sezione di Trieste
Scuola Internazionale Superiore di Studi Avanzati (SISSA)
Via Beirut 2-4, I-34013 Trieste, Italy*

Abstract

It is briefly explained why recent claims about the vanishing of the one-loop effective potential in Matrix theory, thus invalidating the possible agreement with supergravity, do not hold.

In our paper [1], we argued that it is possible for finite- N Matrix theory to reproduce the supergravity amplitude for three-graviton scattering, contrary to what previously claimed in [2, 3]. Our prediction is now fully borne out by the computation of [4], where it is shown that there is perfect agreement between the two theories, numerical coefficients included. In the meantime, two papers [5, 6] came out, based on an effective action approach, arguing that such agreement is impossible. In particular, the two papers disputed our result about the non-vanishing of the relevant one-loop effective potential which we used in discussing the agreement between Matrix theory and supergravity. This short note addresses these objections and show why they do not hold.

Matrix theory is defined by a supersymmetric $SU(N)$ Yang-Mills (YM) field theory in $1 + 0$ dimensions with the classical action

$$S(A_\mu, \Psi) = \int dt \left(-\frac{1}{2} \text{Tr} F^{\mu\nu} F_{\mu\nu} + i \text{Tr} \bar{\Psi} \Gamma \cdot D \Psi \right), \quad (1)$$

where $F_{0k} = \partial_t A_k - i[A_0, A_k]$, $F_{jk} = i[A_j, A_k]$ and $D_0 \Psi = \partial_t \Psi + i[A_0, \Psi]$, $D_k \Psi = i[A_k, \Psi]$. Throughout this note, Greek indices go from 0 to 9 and Latin ones from 1 to 9.

In order to compare the three-graviton scattering in supergravity with finite- N Matrix theory, one has to make a two-loop computation for $SU(3)$ YM in the presence of the following classical external field

$$B_k = \begin{pmatrix} 0 & 0 & 0 \\ 0 & r_2 - r_1 & 0 \\ 0 & 0 & r_3 - r_1 \end{pmatrix}_k, \quad (2)$$

which is solution of the classical equations of motion when the graviton positions $\mathbf{r}_{1,2,3}$ depend linearly on time. We can fix the center of mass by writing $\mathbf{r}_1 + \mathbf{r}_2 + \mathbf{r}_3 = 0$. We assume that $|\mathbf{r}_2 - \mathbf{r}_1|$ and $|\mathbf{r}_3 - \mathbf{r}_1|$ are much larger than $|\mathbf{r}_2 - \mathbf{r}_3|$. Notice that B_k contains an irrelevant decoupled $U(1)$ component that we retain for convenience.

We quantize by setting

$$A_\mu = B_\mu + A_\mu^f, \quad (3)$$

where $B_\mu = (0, B_k)$ and A_μ^f is the quantum (fluctuating) gauge field.

We use the gauge fixing defined by

$$D^\mu(B) A_\mu^f = \partial_t A_0^f + i[B_k, A_k^f]. \quad (4)$$

We have to perform the functional integration

$$e^{-F(B)} = \int [dA_\mu^f][d\Psi][d\bar{C}][dC] e^{-S(B_\mu + A_\mu^f, \Psi) - \int [D^\mu(B) A_\mu^f]^2 - \int \bar{C} D^\mu(B) D_\mu(A) C}. \quad (5)$$

The 2-loop action $F_{(2)}(B)$ is to be compared with the supergravity amplitude for the three-graviton scattering.

We write

$$A_\mu = \begin{pmatrix} a_1 & a_2^\dagger & a_3^\dagger \\ a_2 & Y_{22} & Y_{23} \\ a_3 & Y_{32} & Y_{33} \end{pmatrix}_\mu, \quad (6)$$

where the 2×2 matrix $Y_\mu = B_\mu + Y_\mu^f$ (by a slight abuse of notation, we also denote by B_μ the appropriate 2×2 sub-matrix of (2)); $a_{1,2,3}$ and Y^f are quantum (fluctuating) fields.

For our choice of background (2), $a_{2\mu}, a_{3\mu}$ are the *heavy* modes since their mass terms are proportional to the large distances $|\mathbf{r}_2 - \mathbf{r}_1|$ and $|\mathbf{r}_3 - \mathbf{r}_1|$. Similarly, for the fermions and the ghosts, we denote by $\psi_2, \psi_3, c_2, c_3, \bar{c}_2, \bar{c}_3$ the respective heavy modes, and by $\phi, \mathcal{C}, \bar{\mathcal{C}}$ the remaining *light* fields.

We can then perform the functional integration in two steps. First, we compute at 1-loop

$$e^{-W_{(1)}(Y^f, a_1, \phi, \bar{\mathcal{C}}, \mathcal{C}; B)} = \int [da_2 da_3][d\psi_2 d\psi_3][d\bar{c}_2 d\bar{c}_3][dc_2 dc_3] e^{-S(B_\mu + A_\mu^f, \Psi) - \int [D^\mu(B) A_\mu^f]^2 - \int \bar{C} D^\mu(B) D_\mu(A) C}. \quad (7)$$

To complete the computation one has to integrate over the light modes:

$$e^{-F(B)} = \int [dY^f][da_1][d\phi][d\bar{\mathcal{C}}][d\mathcal{C}] e^{-W_{(1)}(Y^f, a_1, \phi, \bar{\mathcal{C}}, \mathcal{C}; B)}. \quad (8)$$

Therefore, one has to compute

$$W_{(1)} = V_{(1)}(Y^f; B) + \text{terms with } a_1 + \text{derivative terms} + \text{fermionic and ghost terms}, \quad (9)$$

where $V_{(1)}$ is the potential—that is the term with no derivatives—of the field Y^f .

In our paper [1], we have found that $V_{(1)}(Y^f; B) \neq 0$ for generic Y^f . In fact, let us take

$$Y_k^f = \begin{pmatrix} 0 & x \\ x^\dagger & 0 \end{pmatrix}_k, \quad (10)$$

where x_k is the gauge field component in the k direction having the light mass term $|\mathbf{r}_2 - \mathbf{r}_3|$ (it is called $x_k^{\alpha^1}$ in [1]). We obtain that the part of $V_{(1)}$ quadratic in the field $\mathbf{x}$ is

$$V_{(1)}(Y^f; B) = \frac{|\mathbf{R}_1 \cdot \mathbf{x}|^2 - |\mathbf{R}_1|^2 |\mathbf{x}|^2}{R_2 R_3 (R_2 + R_3)}, \quad (11)$$

where $\mathbf{R}_1 = \mathbf{r}_2 - \mathbf{r}_3$ and cyclic.

In the same paper, we have shown that by performing the functional integration over $\mathbf{x}$ and using eq. (11) as its potential, we obtain a term for $F_{(2)}(B)$ which has the same behavior as the supergravity amplitude.

The papers [5] and [6] challenge our result, by claiming that $V_{(1)} = 0$. However, both papers deal with a potential $V_{(1)}$ for field configurations that are not the relevant ones.

In the appendix of ref. [5], it is shown that $V_{(1)}(0; Y) = 0$ for any Y . We have checked this result and found it to be true. This is however irrelevant for the computation of eq. (8), which is the one needed for the comparison with supergravity. In fact, in eq. (8) one must integrate $e^{-W_{(1)}}$ (which contains $V_{(1)}$, see eq. (10)) over the fluctuations Y^f keeping B fixed.

In ref. [6] the computation is done for the case when $\mathbf{R}_1$ and $\mathbf{x}$ are parallel (see eqs. (3) and (34) of ref. [6]). In this case, $V_{(1)} = 0$, which is true, as it can be seen from eq. (11). However, this is just a special choice of $\mathbf{x}$ whereas in eq. (8) one must integrate over all the possible $\mathbf{x}$.

References

- [1] M. Fabbrichesi, G. Ferretti and R. Iengo, *JHEP* 06 (1998) 002 [[hep-th/9806018](#)].
- [2] M. Dine, A. Rajaraman, [hep-th/9710174](#).
- [3] M. Dine, R. Echols and J.P. Gray, [hep-th/9805007](#).
- [4] Y. Okawa and T. Yoneya, [hep-th/9806108](#).
- [5] W. Taylor IV and M. Van Raamsdonk, [hep-th/9806066](#).
- [6] R. Echols and J.P. Gray, [hep-th/9806109](#).